

Glebe Merc
the School

part of 1912

Song of

to 1912

Glebe Merc

Emma Amelia

Carlisle Smith

F = John "Jack"

Sarah Creek

= star of dawn

Pop Randall May

Ed McMillin

Vilate "111111"

Mary's Mother

John M. Munk

Kate Burk & Chas

1/2 sister to Lyman

UTAH INFLUENZA IMMUNIZATION PROJECT

INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at

Hope & Son
1000 Broadway
New York

— 27 —

Tracy

2/3

W. A. R. R.

7/22/21 2122/7

John
Lefkowitz

2-145 41117 10/10

Oct Nov

UTAH INFLUENZA IMMUNIZATION PROJECT

INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at

Ray Mather
Booth Street

How New
1896 27 yr

9990

~~1905~~

The Mathewson

old New
1903

UTAH INFLUENZA IMMUNIZATION PROJECT

INFLUENZA VACCINE REACTION INFORMATION

You have just received influenza vaccine. A small percentage of persons receiving this vaccine may experience one or more of the following symptoms: redness and tenderness at the injection site, fever, (usually 101° or less), chills, nausea, loss of appetite, muscle ache, joint pain, headache, or fatigue. These reactions are usually short lived, lasting less than 48 hours, but local reactions at the place of injection may persist for several days. If you experience severe or prolonged reaction, consult your physician or call your local health department at